ZOOLOGIA 34: e12277
ISSN 1984-4689 (online)
zoologia.pensoft.net

RESEARCH ARTICLE

Flesh flies (Diptera: Sarcophagidae: Sarcophaginae) from the Colombian Guajira biogeographic province, an approach to their ecology and distribution

César Valverde-Castro[1], Eliana Buenaventura[2,3], Juan David Sánchez-Rodríguez[1], Marta Wolff[1]

[1]*Entomology group, Instituto de Biología, Universidad de Antioquia. Calle 67 # 53-108, Laboratory 7-311, Medellín, Colombia.*
[2]*Natural History Museum of Denmark. Universitetsparken 15, DK-2100 Copenhagen, Denmark.*
[3]*National Museum of Natural History, Smithsonian Institution. Washington, DC, United States of America.*
Corresponding author: César Valverde-Castro (cesarvalverdec@gmail.com)

http://zoobank.org/6D556F84-4011-4F8B-A024-E3AD7FE3A4F3

ABSTRACT. Sarcophaginae is the most diverse subfamily of Sarcophagidae with 51 genera and more than 1,800 species and it is widely distributed in the Neotropical Region. Sarcophaginae flies of forest, rural, and urban habitats of the seven departments comprising the Colombian Guajira biogeographic province were surveyed. Van Someren Rydon traps baited with fermented fruit, decomposing fish, and human feces were active for 72 hours with samples collected every 12 hours (day/night) between 2010 and 2014. A total of 14,223 sarcophagines (78% females and 22% males) were collected, which belong to 28 species of nine genera. *Oxyvinia excisa* (Lopes, 1950) is a new record for Colombia. The richest genera were *Oxysarcodexia* Townsend, 1917 and *Peckia* Robineau-Desvoidy, 1830, with eleven and nine species, respectively. Some of the species with the highest affinity to urban habitats were *Oxysarcodexia bakeri* (Aldrich, 1916), *Oxysarcodexia aurata* (Macquart, 1851), and *Oxysarcodexia timida* (Aldrich, 1916); to rural habitats were *Oxysarcodexia diana* (Lopes, 1933), *Ravinia columbiana* (Lopes, 1962), and *Ravinia effrenata* (Walker, 1861); and to forest habitats were *Helicobia rapax* (Walker, 1849), *Oxysarcodexia avuncula* (Lopes, 1933), and *Oxysarcodexia major* Lopes, 1946. We also discuss diet, habitat, and activity periods of the species. Species composition of Sarcophaginae in the Guajira province was very homogenous across the habitats, with the forest as the richest habitat. Scavenger habits of Sarcophaginae are confirmed, as well as their strong association with anthropic environments.

KEY WORDS. Caribbean, Colombia, ecology, Guajira province, Sarcophaginae.

INTRODUCTION

The Neotropical Region comprises three sub-regions (Antillean, Brazilian, and Chacoan), two transition zones (Mexican and South American), seven dominions (Mesoamerican, Pacific, Boreal Brazilian, Southwestern Amazonian, Southeastern Amazonian, Chacoan, and Parana), and 53 provinces (Morrone 2014). The Brazilian sub-region is divided into the Mesoamerican and Pacific dominions. The latter is subdivided into twelve provinces, one of which is the Guajira biogeographic province, which encompasses northern Colombia and northwestern Venezuela (Morrone 2014). Dry forests and shrubs, short trees, scrubs, and cardones (Cactaceae) characterize the Guajira biogeographic province, but it also includes wet forests and swamps maintained by periodic floods (Morrone 2001). This province is also known as the pre-Caribbean arid belt, which is divided into the tropical desert, tropical sub-xerophytic, and hydric zono-biomes (Hernández and Sanchez 1992), and it is nearly 132.000 km^2. According to Holdridge (1967), dry and very dry tropical forests define the Guajira biogeographic province. This biogeographic province is characterized by a selection of endemic species of plants, birds, and beetles (Morrone 2014), although other endemic organisms are present as well, but not yet studied. Some of the few Diptera families reported for this province are Drosophilidae, Ephydridae, Simuliidae (Morrone 2001), and Calliphoridae (Santodomingo et al. 2014). However,

there is no information about Sarcophagidae, known as a widely-distributed family in the Neotropical Region, with species of forensic, medical, and ecological importance (Amorim et al. 2002, Courtney et al. 2009, Carvalho and Mello-Patiu 2008, Ramos and Wolff 2011, Pape and Thompson 2013).

Many flesh fly species seem to be closely associated with humans and domestic animals, and they are found in dung and organic decaying matter (Oliveira et al. 2002). Some species are pathogen vectors of protozoans, helminths, enteropathogenic bacteria, and enteroviruses (Greenberg 1971), and others cause myasis in vertebrates (Dodge 1955, Crump and Pounds 1985, Hagman et al. 2005, Stevens et al. 2006, Bermudez et al. 2010, Mello-Patiu and Luna-Dias 2010). Thus, these flies have medical and veterinary importance. They are also considered a public health problem, mainly because of their eusynanthropic behavior and pathogen vector capacity (Sukontason et al. 2006, Vasconcelos et al. 2015).

Sarcophagidae is the second largest family within Oestroidea, with approximately 171 genera and 3,094 species (Pape et al. 2011). It includes the subfamilies Miltogramminae, Paramacronychiinae, and Sarcophaginae, the latter being the most diverse in the Neotropical Region with around 780 species (Pape 1996). The Sarcophaginae have a large variety of feeding habits, especially in their larval stages. These habits include coprophagy (Pape et al. 2002), parasitism (Dodge 1955, Crump and Pounds 1985, Pape 1994, Stevens et al. 2006, Hagman et al. 2005, Bermúdez et al. 2010, Mello-Patiu and Luna-Días 2010), kleptoparasitism (Spofford and Kurczewski 1989), predation (Pickens 1981, Pape and Dahlem 2010), and necrophagy (Pape 1996, Pape et al. 2004, Barros et al. 2008, Buenaventura et al. 2009, Yepes-Gaurisas et al. 2013). Adult flies feed on nectar, fruit juice, and proteins taken from decomposing matter such as excrements and carrion (Mello-Patiu et al. 2014).

Sarcophagines are pioneers in the entomological succession in human corpses and considered relevant indicators of Post-Mortem Interval (PMI) and other forensic estimations (Pape 1996, Wells et al. 2001, Buenaventura et al. 2009, Buenaventura and Pape 2013). However, their use in criminal investigations is still limited due to: (1) most females and larvae are poorly known, and their taxonomic identification might not be reliable, (2) collection of sarcophagine larvae might not be representative as they spread out up to 10 meters from the cadaver, and (3) knowledge on the ecology, behavior, and distribution is scarce (Jordaens et al. 2013).

The geographic distribution and habitat preference of Neotropical Sarcophagines is still little known but synanthropy studies have provided valuable data in this regard (Ferreira 1979, Linhares 1981, Dias et al. 1984, D'Almeida 1984, Yepes-Gaurisas et al. 2013). In Colombia, records of this fly subfamily are scarce, with the little information being mostly the result of recent studies on forensic entomology (Wolff et al. 2001, Pérez et al. 2005, Martínez et al. 2007, Buenaventura et al. 2009, Barrios and Wolff 2011, Ramos and Wolff 2011, Ramírez-Mora et al. 2012, Amat et al. 2013, Yepes-Gaurisas et al. 2013, Ramos-Pastrana et al. 2014, Mello-Patiu 2016). The catalogue of Sarcophagidae from Colombia (Mello-Patiu 2016) listed 102 species in 26 genera and two subfamilies, of which 96 species belong to Sarcophaginae. In this study, we aimed to contribute to the knowledge on the diversity and ecology of Sarcophaginae, their diet and habitat preferences, as well as their activity periods (day/night) in the Guajira biogeographic province.

MATERIAL AND METHODS

Flies of Sarcofaginae were collected in the western part of the Guajira biogeographic province, which comprises the Colombian departments of La Guajira, Cesar, Magdalena, Atlantico, Bolivar, Sucre, and Cordoba (Fig. 1, Table 1).

Three habitats with differential anthropic intervention were selected in each department. The urban habitat consisted of residential areas in the capital of each department, and was characterized by having drinking water, sewage system, and periodical garbage collection. The rural habitat was an isolated dwelling with septic tanks, farm animals, fishponds, or vegetable crops. The forest habitat was a protected dry forest area.

Six Van Someren Rydon traps placed at 1.5 m height and spaced by 50 m were hung for 72 hours in each habitat on a linear transect of 250 m. Traps were alternately baited with fermented fruit mixture, decomposing fish, and human feces. Specimens were collected every 12 hours (06:00 am and 06:00 pm) and transferred to 96% ethanol. Samples were collected from June 2010 to April 2014 in the three different habitats studied (urban, rural, forest) in each of the seven departments, for a total of 21 localities sampled (Table 1).

Taxonomic identification included only males. Flies were identified using taxonomic keys and descriptions by Mello-Patiu and Pape (2000), Carvalho and Mello-Patiu (2008), Méndez et al. (2008), Buenaventura et al. (2009), Giroux and Wheeler (2009), Pape and Dahlem (2010), Silva and Mello-Patiu (2010), Pinto et al. (2011), Buenaventura and Pape (2013), and Carvalho-Filho et al. (2014). Taxonomic characters of each species were also compared to reference specimens of the Entomological Collection of Universidad de Antioquia, Medellin, Colombia (CEUA). All specimens were deposited in the CEUA (National Record of Biological Collections # 036).

Species preferences for habitat and type of bait were assessed using a Monte Carlo serial analysis (Brower and Kyle 1988), with an input of 30 random matrices. In order to identify the most diverse habitat, we calculated the Shannon-Wiener diversity index (H'). We also tested for statistically significant differences between the abundance and remaining variables using paired Student's t-Tests, considering the normality assumption (Shapiro Wilk). A Correspondence Analysis (CA) was used to estimate the correlation of species vs. habitats and species vs. baits. All analyses were carried out using the statistical program PAST version 3.06 (Hammer et al. 2001).



Figure 1. Map with collection localities in the Guajira province, Colombia.

Table 1. Collection localities in the Colombian Guajira province.

Location in the map	Departments	Municipalities	Location collects	Geographical coordinates	Altitude (msnm)	Habitat
1	La Guajira	Uribia	Vereda Santa Ana	12°19'52.3"N, 71°17'55.5"W	16	Rural
2		Riohacha	Barrio Jorge Pérez	11°32'25.02"N, 72°55'47.45"W	10	Urban
3		Riohacha	Caserío El Abra	11°18'36.69"N, 72°53'55.83"W	44	Forest
4	Magdalena	Santa Marta	SENA Agropecuario	11°11'01.2"N, 74°11'55.1"W	36	Rural
5		Santa Marta	Universidad del Magdalena	11°13'18.36"N, 74°11'10.89"W	20	Urban
6		Santa Marta	Reserva Lalguana Verde	11°10'07.2"N, 74°10'37.4"W	207	Forest
7	Atlántico	Usiacury	Reserva Luriza	10°45'04.99"N, 75°02'33.61"W	125	Rural
8		Barranquilla	Universidad del Atlántico	11°01'07.58"N, 74°52'28.19"W	48	Urban
9		Usiacury	Reserva Luriza	10°45'10.72"N, 75°02'08.59"W	187	Forest
10	Bolívar	Zambrano	Finca La Esmeralda	9°47'38.80"N, 74°50'52.22"W	16	Rural
11		Cartagena	Barrio Las Palmeras	10°23'51.04"N, 75°28'28.29"W	6	Urban
12		Villanueva	Finca El Peligro	10°25'57.59"N, 75°14'52.01"W	302	Forest
13	Cesar	Agustín Codazzi	CORPOICA	10°00'08.4"N, 73°14'54.7"W	113	Rural
14		Agustín Codazzi	Colegio Cooperativo	10°02'31.7"N, 73°14'28.9"W	129	Urban
15		Copey	Vereda Tierras Nuevas	10°13'34.0"N, 73°47'38.9"W	493	Forest
16	Sucre	Rincón del Mar	Cabecera Municipal	9°45'56.69"N, 75°41'00.15"W	2	Rural
17		Sincelejo	Institución Educativa Para Estudiantes Especiales	9°18'49.70"N, 75°23'11.69"W	193	Urban
18		San Onofre	Reserva Sanguaré	9°42'45.09"N, 75°41'00.15"W	16	Forest
19	Córdoba	San Carlos	Vereda Guacharacal	8°33'13.7"N, 75°40'13.6"W	141	Rural
20		Montería	Barrio Santa Fé	8°44'8.7"N, 75°54'26.1"W	47	Urban
21		Montería	Sierra Chiquita	8°43'57"N, 75°54'10.2"W	26	Forest

Names of the genera *Oxysarcodexia* and *Oxyvinia* are abbreviated as "*Oxys.*" and "*Oxyv.*", respectively. Similarly, *Sarcofarhtiopsis* is abbreviated as "*Sarcof.*", *Sarcophaga* as "*Sarcop.*", *Tricharaea* as "*Tr.*", and *Titanogrypa* as "*Ti.*". Genera *Helicobia*, *Peckia*, and *Ravinia* are abbreviated with their inicials.

RESULTS

A total of 14,223 sarcophagines (11,094 females and 3,129 males) were collected, which belong to 28 species of nine genera. *Oxyvinia excisa* (Lopes, 1950) is a new record for Colombia (Table 2). The new records increased to 97 the number of Sarcophaginae species and to 103 the number of flesh flies known to the country.

According to the diversity analysis, the forest was the most homogenous habitat (H' = 2.580), followed by the rural (H' = 2.071) and urban (H' = 2.058) habitats. However, all habitats had regular values as indicated by the Shannon-Wiener diversity index.

Oxysarcodexia bakeri (Aldrich, 1916), *Oxysarcodexia timida* (Aldrich, 1916), *Tricharaea occidua* (Fabricius, 1794), and *Oxysarcodexia conclausa* (Walker, 1861) were the most abundant species in urban environments. The species *Tr. occidua*, *Ravinia columbiana* (Lopes, 1962), *Oxys. conclausa*, *Oxys. bakeri*, *Ravinia effrenata* (Walker, 1861), and *Oxysarcodexia diana* (Lopes, 1933) were the most abundant in rural areas, while *Oxys. conclausa*, *Oxys. timida*, *Tr. occidua*, and *Oxysarcodexia amorosa* (Schiner, 1868) were the most abundant taxa in forest habitats (Table 2).

Table 2. Abundance of Sarcophaginae species (males) collected in the Colombian Guajira province (suc = Sucre, at = Atlántico, mag = Magdalena, lg = La Guajira, ce = Cesar, bl = Bolivar, cor = Cordoba). Species codes in first column are used in the Correspondence Analysis. Taxa are presented by genus, according to their abundance. New record for Colombia are indicated with *.

Genus	Code	Species	Departments	Habitats			Baits			Period of activity		Total
				Urban	Rural	Forest	Fruit	Faeces	Fish	Day	Night	
Oxysarcodexia	1	*Oxys. bakeri* (Aldrich, 1916)	suc, at, mag, lg, ce, bl, cor	263	146	36	81	146	218	344	101	445
	2	*Oxys. conclausa* (Walker, 1861)	suc, at, mag, lg, ce, bl, cor	85	169	136	22	154	214	303	87	390
	3	*Oxys. amorosa* (Schiner, 1868)	suc, at, mag, lg, ce, cor	20	35	70	7	28	90	109	16	125
	4	*Oxys. diana* (Lopes, 1933)	suc, at, mag, ce, bl, cor	29	118	17	5	121	38	152	12	164
	5	*Oxys. sarcinata* Lopes, 1953	suc, at, ce, cor	0	1	15	1	4	11	16	0	16
	6	*Oxys. aurata* (Macquart, 1851)	suc, at, mag, ce, bl, cor	54	29	35	39	12	67	89	29	118
	7	*Oxys. timida* (Aldrich, 1916)	suc, at, mag, ce, bl, cor	228	95	130	50	159	244	322	131	453
	8	*Oxys. angrensis* (Lopes, 1933)	ce, bl, cor	2	0	4	1	0	5	4	2	6
	9	*Oxys. avuncula* (Lopes, 1933)	ce	0	0	9	0	0	9	9	0	9
	10	*Oxys. major* Lopes, 1946	ce	0	0	4	0	0	4	4	0	4
	11	*Oxys. occulta* Lopes, 1946	ce	0	0	2	0	0	2	2	0	2
	12	*P. chrysostoma* (Wiedemann, 1830)	suc, at, mag, lg, ce, bl, cor	41	22	19	12	19	51	69	13	82
	13	*P. pexata* (Wulp, 1895)	mag, lg, ce, bl, cor	1	8	8	2	0	15	15	2	17
	14	*P. lambens* (Wiedemann, 1830)	suc, at, mag, lg, ce, bl, cor	15	10	41	9	0	57	41	25	66
	15	*P. ingens* (Walker, 1849)	at, mag, lg, ce, bl, cor	1	10	17	0	16	12	17	11	29
Peckia	16	*P. gulo* (Fabricius, 1805)	suc	0	0	8	1	0	7	5	3	8
	17	*P. collusor* (Curran & Walley, 1934)	ce	0	0	2	0	2	0	2	0	2
	18	*P. hirsuta* (Hall, 1933)	suc	0	0	1	0	0	1	1	0	1
	19	*P. intermutans* (Walker, 1861)	ce	0	0	1	0	0	1	1	0	1
	20	*P. anguilla* (Curran & Walley, 1934)	ce	0	0	1	0	0	1	0	1	1
Ravinia	21	*R. columbiana* (Lopes, 1962)	suc, at, mag, lg, ce, bl, cor	46	194	57	54	163	80	231	66	297
	22	*R. effrenata* (Walker, 1861)	suc, at, mag, lg, ce, bl, cor	34	144	42	56	109	55	172	48	220
Tricharaea	23	*Tr. occidua* (Fabricius, 1794)	suc, at, mag, lg, ce, bl, cor	120	461	75	44	339	273	505	151	656
Sarcophaga	24	*Sarcop. ruficornis* (Fabricius, 1794)	suc, lg	2	0	0	0	1	1	2	0	2
Titanogrypa	25	*Ti. placida* (Aldrich, 1925)	at, lg, bl	3	0	2	1	0	4	5	0	5
Oxyvinia	26	*Oxyv. excisa* (Lopes, 1950)*	at	0	1	1	0	2	0	2	0	2
Helicobia	27	*H. rapax* (Walker, 1849)	mag	0	0	4	0	0	4	4	0	4
Sarcofahrtiopsis	28	*Sarcof. cuneata* (Townsend, 1935)	lg, cor	0	0	5	5	0	0	5	0	5
Total abundance				944	1443	742	390	1275	1464	2431	698	3129
Number of species				16	15	27	17	15	25	27	16	28



Figure 2. Serial analysis of distribution of species of Sarcophaginae in the Guajira province between habitats (Monte Carlo method with 30 random matrices = 0.878).





Figures 3–4. Correspondence analysis: (3) between the 28 Sarcophaginae species collected in the Guajira province and the three types of habitat (urban, rural, and forest); (4) between species and baits. Species were coded with the numbers from 1 to 28 (see Table 1) to simplify the graphic.

Serial analysis (Fig. 2) showed that *Helicobia rapax* (Walker, 1849), *Oxys. avuncula* (Lopes, 1933), *Oxys. major* (Lopes, 1946), *Oxys. occulta* (Lopes, 1946), *P. anguilla* (Curran & Walley, 1934), *P. collusor* (Curran & Walley, 1934), *P. gulo* (Fabricius, 1805), *P. hirsuta* (Hall, 1933), *P. intermutans* (Walker, 1861), and *Sarcofahrtiopsis cuneata* (Townsend, 1935) are exclusively found in forested areas, while *Sarcophaga ruficornis* (Fabricius, 1794) is found only in urban habitats. In general, we observed that *Oxys. amorosa*, *Oxys. bakeri*, *Oxys. conclausa*, *Oxys. diana*, *Oxys. timida*, *Oxys. aurata* (Macquart, 1851), *P. ingens*, *P. lambens*, *P. chrysostoma* (Wiedemann, 1830), *P. pexata* (Wulp, 1895), *R. columbiana*, *R. effrenata*, and *Tr. occidua* were found in all habitats, making up for 97.8% of the individuals collected (Fig. 2).

Our CA showed association between *H. rapax*, *Oxys. avuncula*, *Oxys. major*, *Oxys. occulta*, *Oxys. sarcinata*, *P. anguilla*, *P. collusor*, *P. gulo*, *P. hirsuta*, *P. intermutans*, and *Sarcof. cuneata* and the forest, and between *Oxys. diana*, *R. columbiana*, *R. effrenata*, and *Tr. occidua* and the rural habitat. *S. ruficornis* was the only taxon associated with the urban habitat, although *Oxys. bakeri*, *Oxys. aurata*, *Oxys. timida*, *P. chrysostoma*, and *Titanogrypa placida* (Aldrich, 1925) showed also slight preference for this habitat. The remaining species did not show a preference for any of the three habitats (Fig. 3).

Paired Student's *t*-Tests showed no significant difference between the habitats, but it revealed differences between the type of baits and the activity periods. This indicates that most of the species were collected during the day with fish and feces bait (Tables 2 and 3).

The most effective bait as measured by species richness was fish with 25 species, followed by fruit with 17, and feces with 15. Fish was also the most effective bait as measured by abundance, attracting 47% of the specimens, followed by feces with 41% and fruit with 12% (Table 1).

Results of the CA between species and baits showed that *Oxyv. excisa* and *P. intermutans* were associated with feces, while *Sarcof. cuneata* showed preference for fruit. Fish was the preferred substrate for most of the species, with the following species showing strong affinity *Oxysarcodexia angrensis* (Lopes, 1933), *Oxys. avuncula*, *Oxys. bakeri*, *Oxys. conclausa*, *Oxys. major*, *Oxys. occulta*, *P. anguilla*, *P. collusor*, *P. gulo*, *P. hirsuta*, *P. lambens*, *P. pexata*, and *Ti. placida*. Other species were collected from three baits without specific affinity to any particular bait (Fig. 4).

According to the CA, all species were active during day except *P. anguilla*, which was collected at night with a single individual (Table 2).

Table 3. Comparison of the abundances of species through paired Student's *t*-Tests between the habitats, baits and period of activity of Sarcophaginae flies in the Colombian Guajira province.

Comparisons		t	(p-value)
Habitats			
Forest	Rural	-1.51385	0.141272
Forest	Urban	-0.393244	0.697117
Urban	Rural	1.15808	0.256615
Baits			
Faeces	Fish	0.139035	0.890418
Faeces	Fruit	2.46099	0.0202824*
Fruit	Fish	-3.14418	0.0039202*
Period of activity			
Day	Night	3.44142	0.0018352*

* Significant difference to 95% confidence intervals.

DISCUSSION

The habitats showed no statistically significant difference in species composition, which indicates that flesh flies are, generally speaking, uniformly distributed throughout the Colombian Guajira province (Suppl. material 1). This region originally comprised mostly dry and very dry tropical forest, located in lowlands below 1,000 m altitude. Due to agricultural expansion (Espinal and Montenegro 1977, Etter et al. 2008) in the lowlands of the region, these forests have been drastically reduced and fragmented, which apparently allows the most adapted species to expand their distribution range.

In terms of dietary habits, Sarcophaginae adult flies showed higher preference for decomposing fish than for feces and fermented fruit. The high frequency of adults visiting the decomposing fish seems to be related to the high fat and protein content in animal tissues, which are needed by females for developing eggs (Pape 1996), as well as for first instar larvae to grow and develop on these substrates (Stevens 2003). Human feces and fruits probably do not provide the necessary amount of nutrients for larval development of Sarcophaginae flies. However, other fly families like Muscidae, Calliphoridae, Drosophilidae, and Stratiomyidae, among others, can develop on these media.

Oxyvinia excisa is here reported for the first time from Colombia. This species has also been found in Brazil and Peru (Pape 1996). In Brazil, specimens of *Oxyv. excisa* were collected in forests using human feces (D'Almeida 1984, Lopes 1973), banana with brown sugar and decomposing "*Coccus comosa*" as baits (possibly *Cocos comosa* [Mart.]) (Lopes 1973). The habits of *Oxyv. excisa* from Peru are unknown. Additionally, the description of *Oxyv. excisa* by Lopes (1950) mentions the saprophagous habit of the larvae. Thus, both the habitat preference and the scavenger habit kown for *O. excisa* are consistent with our results.

The most abundant and generalist species were *Oxys. timida*, *Oxys. bakeri*, *Oxys. conclausa*, *Oxys. diana*, *Oxys. amorosa*, *Oxys. aurata*, *P. chrysostoma*, *R. columbiana*, *R. effrenata*, and *Tr. occidua*, found in all habitats along the Guajira province, and from all baits (Table 2). These species showed necrophagous and coprophagous behaviors. Visits to feces and carrion enable these species to transport pathogenic microorganisms on the adhesive ultrastructures of their legs, which might imply potential risks for public health (Sukontason et al. 2006). Besides this, some species such as *P. lambens* are knowning for cause myasis (Pape 1994, Hall and Wall 1995, Fernandes et al. 2009, Bermudez et al. 2010). As other studies indicated, the saprophagous behavior and preference for decaying substrates of the above metioned species (Oliveira-Costa et al. 2001, Barros et al. 2008), might point to their relevance for forensic estimations in the Guajira province.

Tricharaea occidua was the most abundant and widely distributed species in the Guajira province, with 656 specimens (21% of total specimens collected) (Table 2, Suppl. material 1). This species was also dominant in similar studies in a natural reserve in Buenos Aires, Argentina, where it was found associated with feces (Mariluis et al. 2007, Mulieri et al. 2008). In Brazil, it was the most abundant species in cow's lung tissue (Pereira de Sousa et al. 2015) and the second most abundant in pig carcasses (Barbosa et al. 2009), in buffalo excrements (Marchiori 2014), and the third most abundant in feces in the Archipelago Fernando de Noronha (Couri et al. 2008). In Colombia, Yepes-Gaurisas et al. (2013) recorded *Tr. occidua* feeding on chicken viscera, fish, and human feces. These authors also proposed this species as a potential vector in human disease transmission due to its high abundance, closeness to feces, and strong association with human settlements. However, *Tr. occidua* seems to be able to adapt to different environments and has been recorded in both open habitats and forests (Pereira de Sousa et al. 2015). All above is consistent with the abundance of this species in feces and fish. Our CA showed strong association of *Tr. occidua* with rural habitats, probably because of the abundance of barnyard animal excrements in this environment, as reported by Toyama and Ikeda (1976) in milk farms from Oahu-Hawaii. Our results suggest the importance of this species as a decomposer of carrion and feces and its high dispersion ability in the Neotropical Region.

Oxysarcodexia, with 11 species, was the most diverse and abundant genus in the present study, which confirmed its dominance in the Neotropical Region (Lopes 1973). This genus is considered of importance in forensic entomology studies in Brazil due to its abundance, diversity, and scavenger habits (Dias et al. 1984, Carvalho and Linhares 2001, Oliveira et al. 2002, Barros et al. 2008, Rosa et al. 2009, Barbosa et al. 2009). A study by Oliveira et al. (2002) reported 11 species in Rio de Janeiro, of which *Oxys. thornax* (Walker, 1849) was the most abundant species, while in the present study *Oxys. timida* was the most abundant species of *Oxysarcodexia* with 453 specimens (14.5%). The flesh fly *Oxysarcodexia timida* has been reported as

a PMI indicator (Barbosa et al. 2009, Bitar et al. 2013) and it is strongly associated with urban settlements (Yepes-Gaurisas et al. 2013). Our CA suggests this species as widely distributed in the Colombian Guajira province.

Oxysarcodexia bakeri was the third most abundant species (14.2%). It seems to be a generalist species, since our CA showed no correlation between this species and any particular habitat or bait. Other studies reported coprophagous (Flores and Dale 1995) or necrophagous (Yepes-Gaurisas et al. 2013) habits for this species.

Oxysarcodexia conclausa seems to be common in urban and peri-urban settlements where it showed strong preferences for necrophagous feeding mode (Ramirez-Mora et al. 2012, Yepes-Gaurisas et al. 2013). This is consistent with our results, since the greatest abundance of this species was on fish bait. However, our CA showed no preference of this species for any habitat.

Oxysarcodexia angrensis, *Oxys. amorosa*, *Oxys. occulta*, *Oxys. avuncula*, *Oxys. major*, and *Oxys. sarcinata* were associated with the forest, as confirmed our CA. Similarly, the three latter species were associated with non-disturbed environments and showed necrophagous habits in previous reports (i.e., Yepes-Gaurisas et al. 2013). In addition, *Oxys. avuncula* was found in human corpses (Barros et al. 2008) and pig carrion (Barbosa et al. 2009) in urban and forest areas in Brazil (Dias et al. 1984, Carvalho and Linhares 2001), which suggests its potential as a PMI indicator.

Oxysarcodexia diana, *R. columbiana*, and *R. effrenata* were correlated with rural habitat, which is in agreement with reports by Yepes-Gaurisas et al. (2013), Faria (2013), and Pinilla et al. (2012). Particularly, *Oxys. diana* showed coprophagous and necrophagous habits (D'Almeida 1984, Dias et al. 1984, D'Almeida and Lima 1994, Barbosa et al. 2009). Our CA showed no correlation of *R. columbiana* and *R. effrenata* with any bait, although previous studies reported these species as necrophagous (Yepes-Gaurisas et al. 2013, Faria 2013, Pinilla et al. 2012).

Peckia, with nine species, was the second most abundant genus. Species of this genus have been widely studied due to their forensic importance and diversity (Carvalho and Linhares 2001, Barros et al. 2008, Buenaventura et al. 2009, Barbosa et al. 2009, Rosa et al. 2009, Buenaventura 2013, Buenaventura and Pape 2013, 2015, Yepes-Gaurisas et al. 2013). In Brazil and Colombia, *Peckia* was reported as the second most abundant genus in forensic studies, after *Oxysarcodexia* (Carvalho and Linhares 2001, Barros et al. 2008, Barbosa et al. 2009, Rosa et al. 2009, Yepes-Gaurisas et al. 2013).

We found *P. chrysostoma* as the most abundant species of *Peckia*. This species is widely distributed in the Neotropical Region (D'Almeida 1993, Pape 1996, Buenaventura 2013, Buenaventura and Pape 2013, 2015), and seems to adapt to open habitats and forests (Pereira de Sousa et al. 2015). In Colombia and Brazil, Linhares (1981), D'Almeida (1984), Oliveira et al. (2002), and Yepes-Gaurisas et al. (2013) reported it as eusynanthropic (strong affinity for human-modified habitats) species associated with carrion. In contrast, we found no affinity to any habitat.

Peckia lambens was the second most abundant species of *Peckia* with a wide distribution in all habitats studied in the Guajira province. This species showed necrophagous habit as observed in previous studies (Carvalho and Linhares 2001, Barros et al. 2008, Barbosa et al. 2009, Bitar et al. 2013, Yepes-Gaurisas et al. 2013, Vasconcelos et al. 2015, Pereira de Sousa et al. 2015), one of them including a report in human corpses (Oliveira-Costa et al. 2001). The broad biological spectrum of this species also includes parasitism, as it causes myiasis in birds (Bermúdez et al. 2010), anurans (Hagman et al. 2005), and humans (Queiroz de Leão et al. 1996, Fernandes et al. 2009).

Peckia ingens occurred in all three studied habitats, which is consistent with the study by Yepes-Gaurisas et al. (2013). Our CA showed a correlation of this species with feces and carrion baits. This species might have potential as a PMI indicator in forensic entomology, since previous studies found it as necrophagous on decomposing pig corpses in the Colombian amazon foothills (Ramos and Wolff 2011), and in a forest in Brazil (Carvalho and Linhares 2001).

Peckia pexata was found in all habitats, but it was highly associated with rural and forest habitats. Our CA showed correlation between this species and fish, which indicates necrophagous habits, also confirmed in other studies (Carvalho and Linhares 2001, Barros et al. 2008, Bitar et al. 2013). In Colombia, this species showed low abundance (Yepes-Gaurisas et al. 2013), which is further supported in this study with only 17 specimens collected.

Peckia collusor, *P. anguilla*, *P. hirsuta*, and *P. intermutans* were collected in low numbers in forest habitat. *Peckia collusor* has been collected in decomposing fish (Yepes-Gaurisas et al. 2013) and pig corpses (Carvalho and Linhares 2001, Barbosa et al. 2009, Bitar et al. 2013), which suggests necrophagous habits and some potential as a PMI indicator. In Brazil, *P. intermutans* is a relevant species in forensic entomology due to its scavenger habits (Carvalho and Linhares 2001, Barbosa et al. 2009, Rosa et al. 2009, Oliveira and Vasconcelos 2010, Amorim et al. 2014). According to D'Almeida (1984), *P. intermutans* rejects densely populated areas, which is in agreement with our study as this species was only found in forest habitats.

This is the first ecological approach to the subfamily Sarcophaginae in a specific biogeographic area within Colombia. Our study corroborates the necrophagous habits of some species of this subfamily. Species composition of Sarcophaginae in the Guajira province was very homogenous across the habitats, although the forest showed higher richness.

Historically, Sarcophaginae flies have been associated with anthropic environments, which is consistent with the results presented here. The performance of Sarcophaginae as bacteria carriers and its consequences for public health in this province has not been studied, however the present study highlighted their high dispersion ability and ubiquity in urban habitats. In addition, flesh flies of Sarcophaginae seem to be relevant decomposers of organic matter, thus playing a role as recyclers in terrestrial ecosystems.

ACKNOWLEDGMENTS

We are thankful to the colleagues of the Group of Entomology of Universidad de Antioquia for their support and motivation. We also thank the following institutions for allowing sampling in their facilities: Universidad del Magdalena, Universidad del Atlántico, Secretaria de Salud de Córdoba, Sanguaré reserve, Iguana Verde reserve, Luriza reserve, CORPOICA Codazzi headquarters, SENA agropecuario Santa Marta, Colegio cooperativo, Institución Educativa para Niños Especiales, La Esmeralda farm, and El Peligro farm. C. V-C thanks Mariano Altamiranda, Jaiber Solano, Carlos Villa, Erick Perdomo, Anisbeth Daza, Adriana Santodomingo, Eusebia Betancur, Silvelis De Armas, and Boris Valverde for their help during fieldwork. We thank Universidad de Antioquia (project CODI) for financial support.

LITERATURE CITED

Amat E, Ramírez-Mora MA, Buenaventura E, Gómez-Piñerez LM (2013) Variación temporal de la abundancia en las familias de moscas carroñeras (Diptera, Calyptratae) en un valle andino antropizado de Colombia. Acta Zoológica Mexicana 29: 463–472.

Amorim DS, Silva VC, Balbi MI (2002) Estado do conhecimento dos Diptera neotropicais. In: Costa C, Vanin SA, Lobo JM, Melic A (Eds) Proyecto de Red Iberoamericana de Biogeografia y Entomologia Sistemática PrIBES. Sociedad Entomológica Aragonesa & Cyted, Zaragoza, 2936 pp.

Amorim JA, Souza CM, Thyssen PJ (2014) Molecular characterization of *Peckia (Pattonella) intermutans* (Walker, 1861) (Diptera: Sarcophagidae) based on the partial sequences of the mitochondrial cytochrome oxidase I gene. Journal of Forensic Research 5: 227. https://doi.org/10.4172/2157-7145.1000227

Barbosa RR, Mello-Patiu CA, Mello RP, Queiroz MMC (2009) New records of calyptrate dipterans (Fanniidae, Muscidae and Sarcophagidae) associated with the decomposition of domestic pigs in Brazil. Memórias do Instituto Oswaldo Cruz 104: 923–926. https://doi.org/10.1590/S0074-02762009000600018

Barrios M, Wolff M (2011) Initial study of arthropods succession and pig carrion decomposition in two freshwater ecosystems in the Colombian Andes. Forensic Science International 212: 164–172. https://doi.org/10.1016/j.forsciint.2011.06.008

Barros RM, Mello-Patiu CA, Pujol-Luz JR (2008) Sarcophagidae (Insecta, Diptera) associados à decomposição de carcaças de *Sus scrofa* Linnaeus (Suidae) em área de Cerrado do Distrito Federal, Brasil. Revista Brasilera de Entomología 52: 606–609. https://doi.org/10.1590/S0085-56262008000400011

Bermúdez SE, Buenaventura E, Couri M, Miranda RJ, Herrera JM (2010) Mixed myiasis by *Philornis glaucinis* (Diptera: Muscidae), *Sarcodexia lambens* (Diptera: Sarcophagidae) and *Lucilia eximia* (Diptera: Calliphoridae) over *Ramphocelus dimidiatus* (Avia: Thraupidae) chicks in Panama. Boletín de la Sociedad Entomológica Aragonesa 47: 445–446.

Bitar PDR, Rodrigues TFS, Geiser GC (2013) Ocorrência da família Sarcophagidae (Insecta, Diptera) em carcaças de *Sus scrofa* Linnaeus (Suidae) em Belém-PA: colonização da carcaça e sua relação com o tempo de morte do animal. Revista Brasilera de Criminalística 2: 24–31. https://doi.org/10.15260/rbc.v2i1.44

Brower JC, Kyle KM (1988) Seriation of an original data matrix as applied to palaeoecology. Lethaia 21:79–93. https://doi.org/10.1111/j.1502-3931.1988.tb01756.x

Buenaventura E, Camacho G, García A, Wolff M (2009) Sarcophagidae (Diptera) de importancia forense en Colombia: claves taxonómicas, notas sobre su biología y distribución. Revista Colombiana de Entomología 35: 189–196.

Buenaventura E (2013) Morphology of the first and second instars larvae of *Peckia (Peckia) chrysostoma* (Wiedemann, 1830) (Diptera, Sarcophagidae). Acta Zoológica Mexicana 29: 96–104. https://doi.org/10.21829/azm.2013.291388

Buenaventura E, Pape T (2013) Revision of the new world genus *Peckia* Robineau-Desvoidy (Diptera: Sarcophagidae). Zootaxa 3622: 1–87. https://doi.org/10.11646/zootaxa.3622.1.1

Buenaventura E, Pape T (2015) Phylogeny of the Peckia-genus group: evolution of male genitalia in the major necrophagous guild of Neotropical flesh flies (Diptera: Sarcophagidae). Organisms Diversity and Evolution 15: 301–331. https://doi.org/10.1007/s13127-015-0203-0

Carvalho CJB de, Mello-Patiu CA (2008) Key to the adults of the most common forensic species of Diptera in South America. Revista Brasilera de Entomologia 52: 390–406. https://doi.org/10.1590/S0085-56262008000300012

Carvalho LML, Linhares AX (2001) Seasonality of insect succession and pig carcass decomposition in a natural forest area in southeastern Brazil. Journal Forensic Science 46: 604–608.

Carvalho-Filho FDS, Esposito MC, de Azevedo Silva A (2014) A further new species of *Sarcofahrtiopsis* Hall (Diptera: Sarcophagidae) associated with faeces of the disk-winged bat (*Thyroptera* Spix: Chiroptera) in Brazil and the redescription of the female terminalia of *S. cuneata* (Townsend). Zootaxa 3889: 118–126. https://doi.org/10.11646/zootaxa.3889.1.7

Couri MS, Barros G, Orsini P (2008) Dipterofauna do arquipélogo de Fernando de Noronha (Pernambuco, Brasil). Revista Brasilera de Entomología 52: 588–590. https://doi.org/10.1590/S0085-56262008000400007

Courtney GW, Pape T, Skevington JH, Sinclair BJ (2009) Biodiversity of Diptera. In: Foottit RG, Adler PH (Eds) Insect biodiversity science and society. Wiley-Blackwell, Oxford, 185–222. https://doi.org/10.1002/9781444308211.ch9

Crump ML, Pounds JA (1985) Lethal parasitism of an aposematic anuran (*Atelopus varius*) by *Notochaeta bufonivora* (Diptera: Sarcophagidae). Journal of Parasitology 71: 588–591.

D'Almeida JM (1984) Sinantropia de Sarcophagidae (Diptera) na região metropolitana do Rio de Janeiro. Arquivos da Universidade Federal Rural do Rio de Janeiro 7: 101–110.

D'Almeida JM (1993) Capture of caliptrate flies with different breeding substrates on beaches in Rio de Janeiro, RJ, Brazil. Me-

mórias do Instituto Oswaldo Cruz 88: 215–220. https://doi.org/10.1590/S0074-02761993000200007

D'Almeida JM, Lima SF (1994) Atratividade de diferentes iscas e sua relação com as fases de desenvolvimento ovariano em Calliphoridae e Sarcophagidae (Insecta, Diptera). Revista Brasilera de Entomología 11: 177–186. https://doi.org/10.1590/S0101-81751994000200001

Dias ES, Neves DP, Lopes HS (1984) Estudos sobre a fauna de Sarcophagidae (Diptera) de Belo Horizonte, Minas Gerais. I. Levantamento taxonômico e sinantrópico. Memórias do Instituto Oswaldo Cruz 79: 83–91. https://doi.org/10.1590/S0074-02761984000100010

Dodge HR (1955) Sarcophagid flies parasitic on reptiles (Diptera, Sarcophagidae). Proceedings of the Entomological Society of Washington 57: 183–187.

Espinal LS, Montenegro E (1977) Formaciones vegetales de Colombia. Memoria explicativa sobre el mapa ecológico. Instituto Geográfico "Agustín Codazzi" (IGAC), Bogotá, 201 pp.

Etter A, McAlpine C, Possingham H (2008) Historical patterns and drivers of landscape change in Colombia since 1500: a regionalized spatial approach. Annals of the Association of American Geographers 98: 2–23. https://doi.org/10.1080/000456007017 33911

Faria LS (2013) Entomofauna de potencial interesse forense medico-legal en dois ambientes de área rural de Uberlândia-MG. Uberlândia, Master dissertation, Intituto de Ciências Biomédicas, Univerdidade Federal de Uberlândia, 84 pp. Available online at: https://repositorio.ufu.br/bitstream/123456789/16692/1/Lucas%20Silva.pdf [Accessed: 18/05/2016]

Fernandes F, Pimenta FC, Fernandes FF (2009) First report of human myiasis in Goiás state, Brazil: frequency of different types of myiasis, their various etiological agents, and associated factors. Journal of Parasitology 95: 32–38. https://doi.org/10.1645/GE-1103.1

Ferreira MJM (1979) Sinantropia de dípteros muscóides de Curitiba, Paraná. II Sarcophagidae. Revista Brasilera de Biología 39: 773–781.

Flores V, Dale W (1995) Un estudio sobre ecología de las moscas Sarcophagidae en la costa central peruana. Revista Peruana de Entomología 38: 13–17.

Giroux M, Wheeler TA (2009) Systematics and phylogeny of the subgenus *Sarcophaga* (*Neobellieria*) (Diptera: Sarcophagidae). Annals of the Entomological Society of America 102: 567–587. https://doi.org/10.1111/j.1096-3642.2009.00561.x

Greenberg B (1971) Flies and disease. Ecology, classification and biotic association. Princeton University Press, Princeton, vol. 1.

Hagman M, Pape T, Schulte R (2005) Flesh fly myiasis (Diptera, Sarcophagidae) in Peruvian poison frogs genus *Epipedobates* (Anura, Dendrobatidae). Phyllomedusa 4: 69–73. https://doi.org/10.11606/issn.2316-9079.v4i1p69-73

Hall MJR, Wall R (1995) Myiasis of humans and domestic animals. Advances in Parasitology 35: 257–334. https://doi.org/10.1016/S0065-308X(08)60073-1

Hammer Ø, Harper DAT, Ryan PD (2001) PAST. Paleontological statistics software package for education and data analysis. Palaeontologia Electronica 4: 1–9. http://palaeo-electronica.org/2001_1/past/issue1_01.htm [Accessed: 05/02/2016]

Hernández Y, Sánchez H (1992) Biomas terrestres de Colombia. In: Halffter G (Org.) La diversidad biológica de Iberoamérica. Instituto de Ecología, Xalapa, 154–174.

Holdridge LR (1967) Life zone ecology. Tropical Science Center, San José, 140 pp.

Jordaens K, Gontran S, Richet R, Dupont E, Braet Y, Desmyter S (2013) Identification of forensically important *Sarcophaga* species (Diptera: Sarcophagidae) using the mitochondrial COI gene. International Journal of Legal Medicine 127: 91–504. https://doi.org/10.1007/s00414-012-0767-6

Linhares AX (1981) Synanthropy of Calliphoridae and Sarcophagidae (Diptera) in the city of Campinas, Sao Paulo, Brazil. Revista Brasileira de Entomologia 25: 189–215.

Lopes HS (1950) Novas espécies neotrópicas de *Notochaeta* Aldrich e *Dexosarcophaga* Townsend (Diptera, Sarcophagidae). Revista Brasilera de Biologia 10: 353–364.

Lopes HS (1973) Collecting and rearing sarcophagid flies (Diptera) in Brazil, during 40 years. Anais da Academia Brasileira de Ciências 45: 279–291.

Marchiori C (2014) Species of Diptera of medical, sanitary and veterinary importance collected from Buffalo and Cattle Dung in South Goiás, Brazil. International Journal of Applied Science and Technology 4: 54–57.

Mariluis J, Schnack J, Mulieri P, Torretta J (2007) The Sarcophagidae (Diptera) of the coastline of Buenos Aires city, Argentina. Journal of the Kansas Entomological Society 80: 243–251. https://doi.org/10.2317/0022-8567(2007)80[243:TSDOTC]2.0.CO;2

Martínez E, Duque P, Wolff M (2007) Succession pattern of carrion-feeding insects in Páramo, Colombia. Forensic Science International 166: 182–189. https://doi.org/10.1016/j.forsciint.2006.05.027

Mello-Patiu CA (2016) Family Sarcophagidae. In: Wolff M, Nihei SS, Carvalho CJB (Eds) Catalogue of Diptera of Colombia. Magnolia Press, Auckland, 884–903. https://doi.org/10.11646/zootaxa.4122.1.75

Mello-Patiu CA, Luna-Dias C (2010) Myiasis in the Neotropical amphibian *Hypsiboas beckeri* (Anura: Hylidae) by a new species of *Lepidodexia* (Diptera: Sarcophagidae). Journal of Parasitology 96: 685–688. https://doi.org/10.1645/GE-2423.1

Mello-Patiu CA, Pape T (2000) Definitions of *Dexosarcophaga* Townsend, 1917 and *Sarcofahrtiopsis* Hall, 1933, including two new species and a redescriptions of *Sarcofahrtiopsis cuneata* (Townsend, 1935) (Diptera: Sarcophagidae). Boletín de Entomología Venezolana 15: 181–194.

Mello-Patiu CA, Mariluis JC, Silva KP, Patitucci LD, Mulieri PR (2014) Sarcophagidae. In: Roig-Juñent S, Claps LE, Morrone JJ (Eds) Biodiversidad de artrópodos argentinos. INSUE – UNT ediciones, vol. 4, San Miguel de Tucumán, 475–490.

Méndez J, Mello-Patiu CA, Pape T (2008) New flesh flies (Diptera: Sarcophagidae) from coastal mangroves of Panama with taxonomic notes and keys. Journal of Natural History 42: 249–257. https://doi.org/10.1080/00222930701850216

Morrone J (2001) Biogeografía de América Latina y el Caribe. CYTED, ORCYT-UNESCO, Sociedad Entomológica Aragonesa, Series Manuales y Tesis SEA, Zaragoza, vol. 3, 148 pp.

Morrone J (2014) Biogeographical regionalisation of the Neotropical region. Zootaxa 3782: 1–110. https://doi.org/10.11646/zootaxa.3782.1.1

Mulieri P, Schnack JA, Mariluis JC, Torretta JP (2008) Flesh flies species (Diptera: Sarcophagidae) from a grassland and a woodland in a Nature Reserve of Buenos Aires, Argentina. Revista de Biología Tropical 56: 1287–1294.

Oliveira TC, Vasconcelos SD (2010) Insects (Diptera) associated with cadavers at the Institute of Legal Medicine in Pernambuco, Brazil: implications for forensic entomology. Forensic Science International 198: 97–102. https://doi.org/10.1016/j.forsciint.2010.01.011

Oliveira-Costa J, Mello-Patiu CA, Lopes SM (2001) Dípteros muscóides associados com cadáveres humanos no local da morte, no estado do Rio de Janeiro, Brasil. Boletim do Museu Nacional, Zoologia, 464: 1–7.

Oliveira VC, D'Almeida JM, Paes JM, Sanavria A (2002) Population dynamics of calyptrate Diptera (Muscidae and Sarcophagidae) at the Rio-Zoo Foundation, Rio de Janeiro, RJ, Brazil. Brazilian Journal Biology 62: 191–196. https://doi.org/10.1590/S1519-69842002000200002

Pape T (1994) The world *Blaesoxipha* Loew, 1861 (Diptera: Sarcophagidae). Entomologica Scandinavica 45: 1–247.

Pape T (1996) Catalogue of the Sarcophagidae of the World (Insecta: Diptera). Memoirs on Entomology, International 8: 1–558.

Pape T, Thompson FC (2013) Systema Dipterorum, Nomenclator Status Statistics, version 1.10. Copenhagen, Natural History Museum of Denmark, BioSystematics Database of World Diptera, version 1.5. http://www.diptera.org/FamilyTables.php [Accessed: 10/11/2016]

Pape T, Blagoderov V, Mostovski MB (2011) Order Diptera Linnaeus, 1758. In: Zhang Z-Q (Ed.) Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Magnolia Press, Auckland, 222–229. Available online at: http://www.mapress.com/zootaxa/list/2011/3148.html

Pape T, Dechmann D, Vonhof MJ (2002) A new species of *Sarcofahrtiopsis* Hall (Diptera: Sarcophagidae) living in roosts of Spix's disk winged bat *Thyroptera tricolor* Spix (Chiroptera) in Costa Rica. Journal of Natural History 36: 991–998. https://doi.org/10.1080/00222930110062291

Pape T, Wolff M, Amat E (2004) Los Califóridos, Éstridos, Rinofóridos y Sarcofágidos (Diptera: Calliphoridae, Oestridae, Rhinophoiridae, Sarcophagidae) de Colombia. Biota Colombiana 5: 201–208.

Pape T, Dahlem GA (2010) Sarcophagidae. In: Brown BV, Borkent A, Cumming JM, Wood DM, Woodley NE, Zumbado M (Eds) A Manual of Central American Diptera. NRC Research Press, Ottawa, vol. 2, 1313–1335.

Pereira de Sousa JR, Carvalho-Filho FDS, Esposito MC (2015) Distribution and abundance of necrophagous flies (Diptera: Calliphoridae and Sarcophagidae) in Maranhão, Northeastern Brazil. Journal of Insect Science 15: 1–10. https://doi.org/10.1093/jisesa/iev054

Pérez SP, Duque P, Wolff M (2005) Successional behavior and occurrence matrix of carrion-associated arthropods in the urban area of Medellín, Colombia. Journal of Forensic Science 50: 448–454. https://doi.org/10.1520/JFS2004046

Pickens LG (1981) The life history and predatory efficiency of *Ravinia lherminieri* (Diptera: Sarcophagidae) on the face fly (Diptera: Muscidae). Canadian Entomologist 113: 523–526. https://doi.org/10.4039/Ent113523-6

Pinilla YT, Segura NA, Bello FJ (2012) Synanthropy of Calliphoridae and Sarcophagidae (Diptera) in Bogotá, Colombia. Neotropical Entomology 41: 237–242. https://doi.org/10.1007/s13744-012-0036-x

Pinto E, Vairo KP, Mello-Patiu CA, de Carvalho CJB (2011) Pictorial identification key for species of Sarcophagidae (Diptera) of potential forensic importance in southern Brazil. Revista Brasileira de Entomologia 55: 333–347. https://doi.org/10.1590/S0085-56262011005000033

Queiroz de Leão RN, Fraiha H, Cruz JPN, Tibana R (1996) Miíase uretral por *Sarcodexia lambens* (Wiedemann, 1830) (Diptera: Sarcophagidae). Relato de um caso amazônico. Revista Paraense de Medicina 10: 27–29.

Ramírez-Mora M, Buenaventura E, Gómez-P L, Amat E (2012) Updated checklist and new records of Calyptratae carrion flies (Diptera, Schizophora) from Valle de Aburrá and other localities in Colombia. Entomotropica 27: 27–35.

Ramos Y, Wolff M (2011) Entomofauna cadavérica asociada a cerdos expuestos al sol y sombra, en el pie de monte amazónico colombiano. Momentos de Ciencia 8: 45–54.

Ramos-Pastrana Y, Velasquez-Valencia A, Wolff M (2014) Preliminary study of insects associated to indoor body decay in Colombia. Revista Brasileira de Entomologia 58: 326–332. https://doi.org/10.1590/s0085-56262014005000006

Rosa TA, Babata ML, Souza CM, Sousa D, Mello-Patiu CA (2009) Dipterans of forensic interest in two vegetation profiles of cerrado in Uberlândia, State of Minas Gerais, Brazil. Neotropical Entomology 38: 859–866. https://doi.org/10.1590/S1519-566X2009000600022

Santodomingo-M M, Santodomingo-Santodomingo A, Valverde-C C (2014) Urban blow flies (Diptera: Calliphoridae) of four cities of the Colombian Caribbean Coast. Revista Intrópica 9: 84–91. https://doi.org/10.21676/23897864.1443

Silva KP, Mello-Patiu CA (2010) New species of *Dexosarcophaga* Townsend from Panama with an illustrated key to species of the subgenus *Bezzisca* (Diptera: Sarcophagidae). Journal of Natural History 44: 89–106. https://doi.org/10.1080/00222930903383487

Spofford MG, Kurczewski FE (1989) Nearctic species of Miltogrammini (Diptera: Sarcophagidae) associated with species of Aculeata (Hymenoptera: Vespoidea, Pompiloidea, Sphecoidea, Apoidea). Journal of the Kansas Entomological Society 62: 254–267.

Stevens J (2003) The evolution of myiasis in blowflies (Calliphoridae). International Journal for Parasitology 33: 1105–1113. https://doi.org/10.1016/S0020-7519(03)00136-X

Stevens JR, Wallman JF, Otranto D, Wall R, Pape T (2006) The evolution of myiasis in humans and other animals in the Old and New Worlds (part II): biological and life-history studies. Trends in Parasitology 22: 181–188. https://doi.org/10.1016/j.pt.2006.02.010

Sukontason KL, Bunchu N, Methanitikorn R, Chaiwong T, Kuntalue B, Sukontason K (2006) Ultrastructure of adhesive device in fly in families Calliphoridae, Muscidae and Sarcophagidae, and their implication as mechanical carriers of pathogens. Parasitology Research 98: 477–481. https://doi.org/10.1007/s00436-005-0100-0

Toyama G, Ikeda J (1976) An evaluation of fly breeding and fly parasites at animal farms on Leeward and Central Oahu. Proceedings Hawaiian Entomological Society 2: 353–368.

Vasconcelos SD, Barbosa TM, Oliveira TPB (2015) Diversity of forensically-important dipteran species in different environments in northeastern Brazil, with notes on the attractiveness of animal baits. Florida Entomologist 98: 770–775. https://doi.org/10.1653/024.098.0256

Wells J, Pape T, Sperling FAH (2001) DNA-based identification and molecular systematics of forensically important Sarcophagidae (Diptera). Journal Forensic Science 46: 1098–1102. https://doi.org/10.1520/JFS15105J

Wolff M, Uribe A, Ortiz A, Duque P (2001) A preliminary study of forensic entomology in Medellín, Colombia. Forensic Science International 3058: 1–7.

Yepes-Gaurisas D, Sánchez-Rodríguez JD, Mello-Patiu CA, Wolff M (2013) Synanthropy of Sarcophagidae (Diptera) in La Pintada, Antioquia-Colombia. Revista Biología Tropical 61: 1275–1287. https://doi.org/10.15517/rbt.v61i3.11955

Supplementary material 1

Composition by departments of Sarcophaginae species collected in the Colombian Guajira province

Authors: César Valverde-Castro, Eliana Buenaventura, Juan David Sánchez-Rodríguez, Marta Wolff

Data type: occurence

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/subtbiol.34.12277.suppl1

Submitted: 15 February 2017
Received in revised form: 31 March 2017
Accepted: 11 April 2017
Editorial responsibility: Gabriel L.F. Mejdalani

Author Contributions: CVC and MW designed the experiments; CVC and JDSR conducted the experiments; CVC, JDSR and EB did taxonomic identifications and analyzed the data; CVC, EB and MW wrote the paper.

Competing Interests: The authors have declared that no competing interests exist.